

# The solution of fractional order epidemic model by implicit Adams methods

Ismail Gad, Paolo Novati\*

Department of Mathematics, University of Padova, Italy

March 27, 2015

## Abstract

We consider the numerical solution of the fractional order epidemic model on long time-intervals of a non-fatal disease in a population. Under real-life initial conditions the problem needs to be treated by means of an implicit numerical scheme. Here we consider the use of implicit fractional linear multistep methods of Adams type. Numerical results are presented.

**Keywords:** Fractional linear multistep methods; Fractional order epidemic model; System of nonlinear differential equations.

## 1 Introduction

The problem of spreading of a non-fatal disease in a population which is assumed to have constant size over the period of the epidemic can be formulated in terms of the following first order model (see [1])

$$\begin{cases} x'(t) = -\beta x(t)y(t) \\ y'(t) = \beta x(t)y(t) - \gamma y(t) \\ z'(t) = \gamma y(t) \end{cases} \quad (1)$$

with initial conditions

$$x(t_0) = N_1 \geq 0, \quad y(t_0) = N_2 \geq 0, \quad z(t_0) = N_3 \geq 0, \quad (2)$$

where, at time  $t \geq t_0$ ,  $x(t)$  is the number of susceptible individuals,  $y(t)$  is the number of infected individuals, able to spread the disease by contact with susceptible ones,  $z(t)$  is the number of isolated individuals, who cannot get or transmit the disease for various reasons. Moreover,  $\beta > 0$  is the rate of infection and  $\gamma > 0$  is the rate at which current infective population is isolated. The non-fatality of the model (1) depends on the values of this two parameters. In Figure 1 we show an example of solution of (1) for small initial values.

---

\* E-mail address: novati@math.unipd.it




Figure 1: Plots of numerical solutions for  $x(t)$ ,  $y(t)$ ,  $z(t)$  versus time such that  $N_1 = 20$ ,  $N_2 = 15$ ,  $N_3 = 10$ ,  $\beta = 0.01$  and  $\gamma = 0.02$ .

The fractional order extension of this model have been first studied in [2], where the authors replace the first derivatives in (1) by the Caputo's fractional derivative of order  $0 < \alpha < 1$ , defined by (see e.g. [3])

$$D_{t_0}^{\alpha} f(t) = \frac{1}{\Gamma(1 - \alpha)} \int_{t_0}^t (t - s)^{-\alpha} f'(s) ds,$$

where  $f$  is a given function, and  $\Gamma(\cdot)$  denotes the gamma function. It is known that  $D_{t_0}^{\alpha} f(t) \rightarrow f'(t)$  as  $\alpha \rightarrow 1$ . The main reason that leads to this extension (typically with  $\alpha$  chosen close to 1) is to reduce the error that may arise from neglected parameters or simplifications in the model (1), as for instance the choice of constant rate of infection  $\beta$  and isolation  $\gamma$ . This seems correct in principle since these two parameters may change accordingly with the experience on the spreading of a certain epidemic, that is, on the history of the process. In this view, the use of  $\alpha < 1$  has just the effect of transforming (1) into a model with memory.

Without loss of generality, we assume that the process starts at time  $t_0 = 0$ . Then, using the simplified notation  $f^{(\alpha)}(t) = D_0^{\alpha} f(t)$  we consider the numerical solution of the following fractional system of equations.

$$\begin{cases} x^{(\alpha)}(t) = -\beta x(t)y(t) \\ y^{(\alpha)}(t) = \beta x(t)y(t) - \gamma y(t) \\ z^{(\alpha)}(t) = \gamma y(t) \end{cases} \quad (3)$$

Some of the recent analytic methods for solving nonlinear problems like (??) include the Adomian decomposition method (ADM [?]), homotopy-perturbation

method (HPM [3]), variational iteration method (VIM [?]) and homotopy analysis method (HAM [10,12,13]). They are relatively new approaches to provide an analytical approximate solution (see also [11]) to linear and nonlinear problems and they provide immediate and visible symbolic terms of analytic solutions, as well as numerical approximate solutions to both linear and nonlinear differential equations (see [12–14]). Anyway, such analytic methods are generally effective only for small time intervals.

In this paper we intend to solve (??) by discrete methods for FDEs, since we prefer to avoid restriction on the time interval under investigation. This may be important to forecast the end of the epidemic. Moreover, using real-life values for the initial conditions (2), the problem (3) may become highly stiff and then it will necessary to employ an implicit time-stepping scheme. In this situation, the use of analytic approximation by polynomials, attainable for instance by homotopy analysis method (HAM [2]), seems unreliable. Here, we consider the use of the implicit Fractional Adams methods of order 2, that reduces to the classical trapezoidal rule in the case of  $\alpha = 1$ .

The paper is organized as follows. In Section 2 the stability of the model (3) is investigated. Section 3 is devoted to describe the methods which are used to solve the problem under real-life values for the initial conditions. Numerical results and some implementation details are discussed in Section 4.

## 2 Stability of the model

As mentioned in the introduction, assuming to work with real-life values for the initial conditions, the model (3) can be quite difficult to solve. Indeed, if we consider the Jacobian of the system

$$J = \begin{pmatrix} -\beta y & -\beta x & 0 \\ \beta y & \beta x - \gamma & 0 \\ 0 & \gamma & 0 \end{pmatrix}, \quad (4)$$

its nonzero eigenvalues are given by

$$\lambda_{\pm} = \frac{1}{2}(\beta x - \gamma - \beta y) \pm \frac{1}{2}\sqrt{(\beta x - \gamma - \beta y)^2 - 4\beta\gamma y}.$$

Depending on the values assumed by  $x$  and  $y$  during the process, there is a negative eigenvalue that may be very large. In particular, the stationary point of the component  $y$  of the solution (see Figure 1 and (3)), it is attained for  $\beta x - \gamma = 0$ , where  $y > y_0$ . In this situation

$$\lambda_- = -\frac{\beta y}{2} - \frac{1}{2}\sqrt{\beta^2 y^2 - 4\beta\gamma y}$$

In most of papers that considers the numerical solution of this problem, the initial values and the parameters are taken such that  $\beta y \approx 1$  or more generally very small. In this situation the problem can be solved efficiently by an explicit

scheme since the eigenvalues remains close to the origin. Using more realistic (large) values for  $N_i$ , we clearly have that around the stationary point

$$\lambda_- \approx -\beta y < -\beta N_2$$

so that the problem needs to be solved by an implicit scheme.

### 3 Description of the numerical methods

In this section we present the methods used in the paper, that, depending on the initial conditions, can be used to solve the epidemic model without restrictions on the time length. Consider the general initial value problem

$$D^\alpha y(t) = f(t, y(t)), \quad y(t_0) = y_0, \quad 0 < \alpha \leq 1, \quad t_0 < t \leq T. \quad (5)$$

In a discrete numerical method the time interval  $[t_0, T]$  is replaced by a discrete set of points  $t_j = t_0 + jh$ ,  $h = \frac{T-t_0}{N}$ ,  $j = 0, 1, \dots, N$ , so that the solution is approximated by a sequence  $\{y_j\}_{j=0,1,\dots,N}$  such that  $y_j \approx y(t_j)$ .

The exact solution of (5) can be written in terms of a Volterra integral equation of the second kind with a weakly singular kernel,

$$y(t) = y(t_0) + \frac{1}{\Gamma(\alpha)} \int_{t_0}^t (t-s)^{\alpha-1} f(s, y(s)) ds \quad (6)$$

so that we can write

$$y(t_{k+1}) = y(t_0) + \frac{1}{\Gamma(\alpha)} \sum_{j=0}^k \int_{t_j}^{t_{j+1}} (t_{k+1}-s)^{\alpha-1} f(s, y(s)) ds$$

In this way we can construct a numerical method by considering the product trapezoidal rule in each sub-interval, obtaining the approximation

$$\frac{1}{\Gamma(\alpha)} \sum_{j=0}^k \int_{t_j}^{t_{j+1}} (t_{k+1}-s)^{\alpha-1} f(s, y(s)) ds \approx \frac{h^\alpha}{\Gamma(\alpha+2)} \sum_{j=0}^{k+1} a_{j,k+1} f(t_j, y(t_j)) \quad (7)$$

where

$$a_{j,k+1} = \begin{cases} k^{\alpha+1} - (k-\alpha)(k+1)^\alpha, & \text{if } j=0, \\ (k-j+2)^{\alpha+1} + (k-j)^{\alpha+1} - 2(k-j+1)^{\alpha+1}, & \text{if } 1 \leq j \leq k, \\ 1, & \text{if } j=k+1. \end{cases}$$

The approximation (7) naturally leads to the implicit formula

$$y_{k+1} = y_0 + \frac{h^\alpha}{\Gamma(\alpha+2)} \left( \sum_{j=0}^k a_{j,k+1} f(t_j, y_j) + f(t_{k+1}, y_{k+1}) \right), \quad (8)$$

that generalizes to the fractional case the standard one-step Adams-Multon method of order 2. In order to compute  $y_{k+1}$  from (8) we can replace  $y_{k+1}$  by a predictor approximation  $y_{k+1}^P$ , and then use a corrector iteration

$$\begin{aligned} y_{k+1}^{(i)} &= y_0 + \frac{h^\alpha}{\Gamma(\alpha + 2)} \left( \sum_{j=0}^k a_{j,k+1} f(t_j, y_j) + f(t_{k+1}, y_{k+1}^{(i-1)}) \right), \quad (9) \\ y_{k+1}^{(0)} &: = y_{k+1}^P, \quad i = 1, \dots, q. \end{aligned}$$

The remaining problem is the determination of the predictor formula that we require to calculate the value  $y_{k+1}^P$ . Here, we use the generalized one-step Adams-Bashforth method (that is, the Fractional Euler method) that consists in replacing the integral on the right-hand side of Equation (6) by the product rectangle rule

$$\int_{t_0}^{t_{k+1}} (t_{k+1} - s)^{\alpha-1} f(s, y(s)) ds \approx \frac{h^\alpha}{\alpha} \sum_{j=0}^k b_{j,k+1} f(t_j, y(t_j)),$$

where now

$$b_{j,k+1} = (k + 1 - j)^\alpha - (k - j)^\alpha.$$

Thus, the predictor  $y_{k+1}^P$  is determined by the explicit recursion

$$y_{k+1}^P = y_0 + \frac{h^\alpha}{\Gamma(\alpha + 1)} \sum_{j=0}^k b_{j,k+1} f(t_j, y_j). \quad (10)$$

The method (9) has been considered and discussed in [15].

Besides the predictor corrector implementation described by (9) and (10), in this work we also consider the fully implicit implementation of (8), in which  $y_{k+1}$  is computed by solving with Newton iteration the nonlinear system

$$\Phi(y_{k+1}) - g_k = 0$$

where

$$\begin{aligned} \Phi(y_{k+1}) &: = y_{k+1} - \frac{h^\alpha}{\Gamma(\alpha + 2)} f(t_{k+1}, y_{k+1}) \\ g_k &: = y_0 + \frac{h^\alpha}{\Gamma(\alpha + 2)} \left( \sum_{j=0}^k a_{j,k+1} f(t_j, y_j) \right) \end{aligned}$$

and for implementation, we need to work with the Jacobian (4) of the system.

Finally, we recall the most significant results on the error analysis of the method (8). The first result is based on smoothness assumptions on  $D^\alpha y$ ; the second one on smoothness assumptions on the solution  $y$ .

**Theorem 1.** *reference* Let  $\alpha > 0$  and assume that  $D^\alpha y \in C^2[t_0, T]$  for some suitable  $T$ . Then,

$$\max_{0 \leq j \leq N} |y(t_j) - y_j| = \begin{cases} O(h^2) & \text{if } \alpha \geq 1, \\ O(h^{1+\alpha}) & \text{if } \alpha < 1. \end{cases}$$

**Theorem 2.** *reference* Let  $0 < \alpha < 1$  and assume that  $y \in C^2[t_0, T]$  for some suitable  $T$ . Then, for  $1 \leq j \leq N$  we have

$$|y(t_j) - y_j| \leq C t_j^{\alpha-1} \times \begin{cases} h^{1+\alpha} & \text{if } 0 < \alpha < \frac{1}{2}, \\ h^{2-\alpha} & \text{if } \frac{1}{2} \leq \alpha < 1, \end{cases}$$

where  $C$  is a constant independent of  $j$  and  $h$ .

Thus for all choices of  $\alpha > 0$  the described method gives a convergence order of at least one if either  $y$  or  $D^\alpha y$  is at least two times continuous differentiable on  $[t_0, T]$ .

## 4 Numerical results and discussion

In this section we present some numerical experiments for the model (??), using different values for the initial conditions. We consider relatively long time intervals and different values of  $\alpha$ . In particular we work with  $\alpha = 1, 0.99, 0.95$  represented respectively by a solid, dashed and solid line in each picture. We always take  $\beta = 0.01$  and  $\gamma = 0.02$ .

In Figure 2, for small initial conditions, we consider the Generalized Euler Method (GEM) given by

$$y(t_{k+1}) = y(t_k) + \frac{h^\alpha}{\Gamma(\alpha + 1)} f(t_k, y(t_k)),$$

that of course is an explicit code. This method has been used in [20] for fractional order models of HTLV infection, in [21] to study the HIV during the primary infection and in [22] for the problem of the population dynamics of the human immunodeficiency type 1 virus (HIV-1). More details on this approach can be found in [23] In Figure ?? we plot the numerical solution given by the predictor corrector formula (9)-(10) for higher values of the initial conditions.  $N_1 = 10^4$ ,  $N_2 = 10^3$  and  $N_3 = 10$ . We take  $h = 0.01$  and apply only one corrector iteration. For this problem GEM produces unstable solutions unless we take  $h = \text{explain please}$  Using the same timestep  $h = 0.01$ , the predictor corrector formula shows a certain instability at the beginning of the process when  $N_2 \geq 10^4$ , as shown in Figure 4. This phenomenon seems independent of the number of corrector iterations. The fully implicit implementation of (8) has been used to solve the problem with  $N_1 = 10^5$ ,  $N_2 = 10^4$  and  $N_3 = 10^3$  (timestep  $h = 0.01$  correct??). The results are reported in Figure 5.




Figure 2: Numerical solutions by GEM for  $N_1 = 499$ ,  $N_2 = 10$  and  $N_3 = 1$  (see [9]). The timestep is  $h = 0.001$

## 5 Conclusion

In this paper we discussed numerical methods to obtain the solution of fractional epidemic model (3) over a long time period where HAM [2, 8] is not effective. Increasing the initial conditions the problem becomes difficult to solve. In particular, under realistic values of the initial conditions  $N_1, N_2 \geq 10^4$  with  $\beta = 0.01$ ,  $\gamma = 0.02$ , only the fully implicit implementation of formula (8) is able to provide accurate solution with a reasonable choice of the timestep.

## References

- [1] D.W. Jordan, P. Smith, *Nonlinear Ordinary Differential Equations, third ed.*, Oxford University Press, (1999).
- [2] S. Z. Rida, A.S. Abdel Rady, A.A.M. Arafa, M. Khalil, Approximate analytical solution of the fractional epidemic model, *IJAMR*, **1**, (2012), 17-29.
- [3] I. Podlubny, *Fractional Differential Equations*, Academic Press, San Diego-New York-London, (1999).
- [4] J. Biazar, Solution of the epidemic model by Adomian decomposition method, *Applied Mathematics and Computation*, **173**, (2006), 1101-1106.
- [5] M. Rafei, D.D. Ganji, H. Daniali, Solution of the epidemic model by homotopy perturbation method, *Applied Mathematics and Computation*, **187**, (2007), 1056-1062.




Figure 3: Numerical solutions by the predictor corrector formula (9)-(10) with initial conditions  $N_1 = 10^4$ ,  $N_2 = 10^3$  and  $N_3 = 10$ . The timestep is  $h = 0.01$ .

- [6] M. Rafei, D.D. Ganji, H. Daniali, Variational iteration method for solving the epidemic model and the prey and predator problem, *Applied Mathematics and Computation*, **186**, (2007), 1701-1709.
- [7] Abdul-Monim Batiha, Belal Batiha, A New Method for solving epidemic model, *Australian Journal of Basic and Applied Sciences*, **5**(12), (2011), 3122-3126.
- [8] O. A. Arqub, A. El-Ajou, Solution of the fractional epidemic model by homotopy analysis method, *Journal of King Saud University - Science*, **25**(1), (2013), 73-81.
- [9] Fadi Awawdeh , A. Adawi, Z. Mustafa, Solutions of the SIR models of epidemics using HAM, *Chaos, Solitons and Fractals*, **42**(5), (2009), 3047-3052.
- [10] M. Zurigat, S. Momani, A. Alawneh, Analytical approximate solutions of systems of fractional algebraic differential equations by homotopy analysis method, *Comput. Math. Appl*, **59**(3), (2010), 1227-1235.
- [11] M. Zurigat, S. Momani, A. Alawneh, The homotopy analysis method for handling systems of fractional differential equations, *Appl. Math. Model*, **34**(1), (2010), 24-35.
- [12] M. Merdan, A. Gökdoğan, A.T. Yildirim, On the numerical solution of the model for HIV infection of  $CD4^+$ T-cells, *Comput. Math. Appl.*, **62**(1), (2011), 118-123.






Figure 4: Numerical solutions by the predictor corrector formula (9)-(10) with initial conditions  $N_1 = 10^4$ ,  $N_2 = 10^4$ ,  $N_3 = 10^2$ , and timestep  $h = 0.01$ . The method is implemented with %%%%% ??? %%%%% number of corrector iterations.

- [13] A.M.A. El-Sayed, S.Z. Rida, A.A.M. Arafa, On the solutions of time-fractional bacterial chemotaxis in a diffusion gradient chamber, *Int. J. Nonlinear Sci.*, **7**(4), (2009), 485-492.
- [14] I. Hashim, O. Abdulaziz, S. Momani, Homotopy analysis method for fractional IVPs, *Commun. Nonlinear Sci. Numer. Simul.*, **14**(3), (2009), 674-684.
- [15] K. Diethelm and A. D. Freed, The FracPECE subroutine for the numerical solution of differential equations of fractional order, in: S. Heinzl, T. Plesser (eds.), *Forschung und wissenschaftliches Rechnen 1998*, (1999a), 57-71.
- [16] K. Diethelm and A. D. Freed, On the solution of nonlinear fractional differential equations used in the modeling of viscoplasticity, in F. Keil, W. Mackens, H. Voß and W. J.(eds), *Scientific Computing in Chemical Engineering II: Computational Fluid Dynamics, Reaction Engineering, and Molecular Properties* (Springer, Heidelberg), (1999b), 217-224.
- [17] K. Diethelm and N. J. Ford, Numerical solution of the Bagley-Torvik equation, *BIT*, **42**(3), (2002), 490-507.
- [18] K. Diethelm, N. J. Ford, and A. D. Freed, Detailed error analysis for a fractional Adams method, *Numer. Algorithms*, **36**(1), (2004), 31-52.




Figure 5: Numerical solutions by (8) with Newton iteration for  $x(t)$ ,  $y(t)$ ,  $z(t)$  versus time such that  $N_1 = 10^5$ ,  $N_2 = 10^4$  and  $N_3 = 10^3$ .

- [19] K. Diethelm, N. J. Ford, and A. D. Freed, A predictor-corrector approach for the numerical solution of fractional differential equations, *Nonlinear Dynamics*, **29**(1-4), (2002), 3-22.
- [20] A.A.M. Arfa, S.Z. Rida, M. Khalil, Fractional order model of human T-cell lymphotropic virus I (HTLV-I) infection of CD4<sup>+</sup>T-cells, *Adv. Stud. Biol.*, **3**(7), (2011), 347-353.
- [21] A.A.M. Arfa, S.Z. Rida, M. Khalil, Fractional modeling dynamics of HIV and CD4<sup>+</sup>T-cells during primary infection, *Nonlinear Biomed. Phys.*, **6**(1), (2012), 1-7.
- [22] A.A.M. Arfa, S.Z. Rida, M. Khalil, The effect of anti-viral drug treatment of human immunodeficiency virus type 1 (HIV-1) described by a fractional order model, *Appl. Math. Modelling*, **37**(4), (2013), 2189-2196.
- [23] Z. Odibat, S. Moamni, An algorithm for the numerical solution of differential equations of fractional order, *J. Appl. Math and Informatics.*, **26**(1 - 2), (2008), 15-27.
- [24] Ch. Lubich, Discretized fractional calculus, *SIAM J. Math. Anal.*, **17**(3), (1986), 704-719.
- [25] E. Hairer, C. Lubich, and M. Schlichte. Fast numerical solution of weakly singular Volterra integral equations, *J. Comput. Appl. Math.*, **23**(1), (1988), 87-98.